


A comparison of faecal glucocorticoid metabolite concentration and gut microbiota diversity in bonobos (*Pan paniscus*)

Alexana J. Hickmott^{1,2,3,*}, Klaree J. Boose¹, Monica L. Wakefield⁴, Colin M. Brand^{5,6}, J. Josh Snodgrass¹, Nelson Ting^{1,7} and Frances J. White¹



Abstract

Sex, age, diet, stress and social environment have all been shown to influence the gut microbiota. In several mammals, including humans, increased stress is related to decreasing gut microbial diversity and may differentially impact specific taxa. Recent evidence from gorillas shows faecal glucocorticoid metabolite concentration (FGMC) did not significantly explain gut microbial diversity, but it was significantly associated with the abundance of the family Anaerolineaceae. These patterns have yet to be examined in other primates, like bonobos (*Pan paniscus*). We compared FGMC to 16S rRNA amplicons for 202 bonobo faecal samples collected across 5 months to evaluate the impact of stress, measured with FGMC, on the gut microbiota. Alpha diversity measures (Chao's and Shannon's indexes) were not significantly related to FGMC. FGMC explained 0.80% of the variation in beta diversity for Jensen–Shannon and 1.2% for weighted UniFrac but was not significant for unweighted UniFrac. We found that genus *SHD-231*, a member of the family Anaerolinaceae had a significant positive relationship with FGMC. These results suggest that bonobos are relatively similar to gorillas in alpha diversity and family Anaerolinaceae responses to FGMC, but different from gorillas in beta diversity. Members of the family Anaerolinaceae may be differentially affected by FGMC across great apes. FGMC appears to be context dependent and may be species-specific for alpha and beta diversity but this study provides an example of consistent change in two African apes. Thus, the relationship between physiological stress and the gut microbiome may be difficult to predict, even among closely related species.

INTRODUCTION

The gastrointestinal tract (gut) microbiota play essential roles in host nutrition and health across mammals [1–3]. Perturbations and dysbiosis to gut microbial communities have been linked to diseases like obesity, diabetes, irritable bowel disorders and certain cancers in humans [4–8]. At the same time, physiological stress, hereafter referred to as stress, has been found to have a negative effect on the gastrointestinal tract and associated microbiota [9]. The definition of stress is a threat to homeostasis, which can be acute or chronic [10, 11]. Chronic stress is stress that is experienced over a longer time frame (e.g. days to years), while acute stress often only lasts for several minutes to hours [10, 11]. As used in the human and non-human primate (NHP) literature, stress often refers to psychosocial stress or stress caused by a response to social stimuli that disrupts the normal physiological equilibrium. Stress will refer to chronic stress, including psychosocial stress and other types of stress like dietary stress. However, determining the cause of stress can be difficult, and a very fine level of both behavioural and biomarker data is needed to determine the source of stress. Stress has been linked to dysbiosis, or a disruption to the homeostasis of the microbial community, in the gut microbiota in several mammalian taxa, including humans [12–16, 17], and has also been implicated in mediating the communication between a host and their commensal gut microbiota [18, 19]. The gastrointestinal tract and its associated microbiota play essential roles in host nutrition and health across mammals [1–3].

Perturbations and dysbiosis to gut microbial communities have been linked to diseases like obesity, diabetes, irritable bowel disorders and certain cancers in humans [4–8]. At the same time, stress has been found to have a negative effect on the gastrointestinal tract and associated microbiota for humans in the clinical setting [9]. Stress hormones have been proposed as a mechanism for communication along the gut–brain axis [14]. The gut–brain axis is a bidirectional signalling pathway between the brain and the gut that is potentially mediated by gut microbiota [20]. For example, in humans, stress has been linked to a decrease in the number of species found in the gut microbiota [9]. Additionally, evidence in humans suggests a link between stress, the gut microbiota and immune system function [21]. This evidence from humans is also recapitulated in laboratory models [10, 22–25]. In germ-free rats, the lack of a gut microbial community increases a rat's behavioural and endocrine stress response [22]. Other lab studies in mice have linked depression symptoms, anorexia and cancer to increased stress levels and disruptions in the murine gut microbiota [23–26].

Stress and the gut microbiota has primarily been examined in laboratory models, with very few studies looking at wild-living mammals [14]. In wild-living eastern grey squirrels, physiological indicators of stress were measured in conjunction with gut microbial composition, physiological stress better explained gut microbiota diversity, or metrics that summarize how abundant and the types of bacterial and archaeal species that are in a microbial community, compared to environmental factors [27]. A recent study on pangolins found those raised in captivity had higher microbial diversity associated with lower stress than those rescued from the wildlife trade [28]. In elephants, a stressful event such as translocation appears to induce alterations to the microbiome especially in taxa Planococcaceae, Clostridiaceae, Spirochaetaceae and Bacteroidia increased after the elephants shifted to living in a captive environment [29]. Research into glucocorticoid concentration in rhinos found that about 10% of taxa were related to glucocorticoid concentrations with Aerococcaceae, Atopostipes, Carnobacteriaceae and Solobacterium differentially increased [30]. Therefore, there seems to be a general mammalian pattern that suggests higher stress is associated with increases in specific taxa. However, this pattern has not been well-studied across mammals, and the specific taxa that increase seem to vary widely across mammalian groups.

Amongst NHPs, great apes exhibit similar psychosocial and ecological pressures making them an excellent model to understand how stress and the gut microbiome co-vary [17, 31–34]. The mammalian pattern of higher stress being associated with a more disrupted gut microbiome [27–29, 34] was not found in a study of wild-living western lowland gorillas (*Gorilla gorilla gorilla*) [35]. Vlčková *et al.* [35] found no relationship between alpha and beta diversity measures and proximate stress measures in this

Received 07 January 2022; Accepted 28 June 2022; Published 12 August 2022

Author affiliations: ¹Department of Anthropology, University of Oregon, Eugene, OR 97403, USA; ²Texas Biomedical Research Institute, San Antonio, TX 78227, USA; ³Southwest National Primate Research Center, San Antonio, TX, USA; ⁴Sociology, Anthropology, and Philosophy, Northern Kentucky University, Highland Heights, KY 41099, USA; ⁵Department of Epidemiology and Biostatistics, University of California, San Francisco, USA; ⁶Bakar Computational Health Sciences Institute, University of California, San Francisco, USA; ⁷Institute of Ecology and Evolution, University of Oregon, Eugene, OR 97403, USA.

*Correspondence: Alexana J. Hickmott, ahickmott@txbiomed.org

Keywords: bonobo; cortisol; faecal glucocorticoid metabolites; microbiota; stress.

Abbreviations: ANCOM, analysis of composition of microbiomes; ANOVA, analysis of variance; ASV, amplicon sequence variants; DRC, Democratic Republic of Congo; FGMC, faecal glucocorticoid metabolite concentration; NHP, non-human primate; PERMANOVA, permutational multivariate analysis of variance.

Repositories: This Targeted Locus Study project has been deposited at DDBJ/EMBL/GenBank under the accession KFUK00000000. The version described in this paper is the first version, KFUK01000000, see also: https://www.ncbi.nlm.nih.gov/biosample?LinkName=bioproject_biosample&from_uid=794787, SUBID = SUB10902613, BioProject = PRJNA794787, BioSample = SAMN24657338; Accession = KFUK00000000, and Organism = gut metagenome to find the data.

Nineteen figures are available with the online version of this article.

species. Vlčková and colleagues also found a positive correlation between proximate measures of stress and relative abundance of three different gut microbial taxa (family Anaerolineaceae, genus *Clostridium* cluster XIVb and genus *Oscillibacter*), suggesting that stress is associated with increases in certain types of bacteria within the gastrointestinal tract [35]. These results from western lowland gorillas indicate stress may not have as significant an effect on NHP gut microbial diversity but follows the mammalian trend of having specific taxa differentially abundant with increasing stress. Bonobos (*Pan paniscus*) and gorillas face similar social stressors in that they both are group-living great apes experiencing affiliative and aggressive interactions with conspecifics [36]. Both also tend to live in mixed age and mixed-sex groups, though in western lowland gorillas, there is typically only one male [37–42]. Whether all NHPs have the same taxa that increase with increasing stress and whether they show similar stability in how the diversity of the gut microbiota remains stable has yet to be determined. Additionally, the amounts and effects of stress hormones vary across different groups of mammals and NHPs and may be different in closely related taxa, like gorillas and bonobos.

Stress hormones, specifically glucocorticoids such as cortisol, are considered the principal chemical compounds involved in the stress system of mammals, including NHPs [18, 43]. Glucocorticoids are a type of steroid hormone that fall into the class of corticosteroids, and cortisol, the primary mammalian stress hormone, is one type of glucocorticoid involved in the stress response [18]. Once the adrenal gland excretes cortisol, it is broken down and metabolized [44]. These metabolites are then excreted through saliva, faeces and urine and can be incorporated into tissues like nails and hair [45–48]. The concentration of these metabolites in faeces are referred to as faecal glucocorticoid metabolite concentration (FGMC), and are known to be related to chronic stress rather than acute stress [49, 50]. Therefore, FGMC are capturing the stress an individual experienced in the preceding 48 h before a faecal sample was collected [51]. It is of note that sometimes low stress can be an indicator of suppression of the stress response as well as low stress. The relationship between the gut microbiota and hormonal systems has far-reaching implications for host physiology [9–11, 52]. Nevertheless, how a host's gut microbiota responds to various stress-based fluctuations during short-term variation in stress remains to be examined in many NHPs, like bonobos.

Stress has been studied in wild-living bonobos as it relates to sociality and socio-sexual behaviour [31, 45, 53, 54]. Bonobos are female bonded, male philopatric and exhibit a fission-fusion social system, and this social structure may contribute to the sex-based patterns in FGMC [38, 39, 55, 56]. Sex differences in bonobo stress have been quantified in inter-group encounters with both females and males exhibiting higher cortisol during intergroup encounters but with males having overall higher levels of cortisol [53]. Additionally, captive bonobos exhibited a similar pattern where the single male had overall higher cortisol levels than the five other females [48]. These physiological patterns and social structure more likely emulate that of the *Pan-Homo* common ancestor as compared to gorillas making bonobos an ideal model in which to study stress and the microbiome.

We aim to use bonobos as a model to test patterns of NHP stress and gut microbiota. We predict that bonobo gut microbiota will exhibit a pattern similar to what was found in western lowland gorillas because gorillas and bonobos live in a similar social and ecological environment. Gorillas and bonobos have been hypothesized to use similar resources, like terrestrial herbaceous vegetation, perhaps buffering dietary stress for both great ape populations, unlike chimpanzees. Bonobos and gorillas may have similar gut microbiota because they are both great ape species or because of similar environments [54–58]. Bonobos and gorillas diverged 8–19 million years ago and, therefore, may exhibit differences due to phylogenetic differences [59, 60]. However, the gut microbiota may alternatively have similar responses to stress due to these ecological similarities between bonobos and gorillas. If they are very different, then it suggests that ecological environments are not the important factor in a gut microbiota's response to FGMC. We predict that alpha diversity, or within individual diversity, will not be significantly related to FGMC. We predict that beta diversity, or between individual diversity, will not be significantly related to FGMC. We expect to find several taxa's abundance, specifically family Anaerolineaceae, genus *Clostridium* cluster XIVb and genus *Oscillibacter*, to be significantly explained by FGMC.

METHODS

Study site and sample collection

The research site was the Iyema field camp, located in the Lomako-Yokokala Faunal Reserve just north of the Lomako river at (00°55' North, 21°06' East) in the Democratic Republic of Congo (DRC). The site was mainly covered by primary forest in *terra firma* soil with swamps [54, 57, 58]. We followed bonobos to their night nests for data collection as part of the African Wildlife Foundation's habituation efforts from June 2017 to October 2017. Night nest locations were marked, and each nesting site was revisited the following day. We identified each bonobo as it exited the nest and collected approximately 5 g of faecal sample into 50 ml tubes with 10 ml of RNAlater for each individual in the nesting party [58]. While there is some debate about the effectiveness of different sample preservation methods for examining gut microbiomes [61–67], there is no clear present consensus. We used RNAlater here due to field site remoteness and downstream host genetic analyses.

The samples were stored in a cool, dry place from June–October 2017 until shipped to the Ting Laboratory at the University of Oregon. They were then stored in a minus 20 °C freezer until extraction. The remainder of each faecal sample not collected in RNAlater was brought back to camp, dried using a camp stove and placed into bags with desiccant for FGMC analysis. Thus, for

each faecal sample, we can obtain data on gut microbiota composition and diversity and FGMC. We collected 218 paired faecal samples.

DATA COLLECTION

ELISA assays – FGMC

To evaluate FGMC, we analysed 218 dried faecal samples in the Global Health Biomarker Laboratory at the University of Oregon using ELISA assays to quantify cortisol as a measure of FGMC. We used the Arbour Assay's DetectX Cortisol Enzyme Immunoassay Kit (Arbour Assay's DetectX cat. no. K003-H5W), as it is designed to be used on dried faecal samples and was previously validated for bonobos [68, 69]. We included known controls provided for Cincinnati Zoo bonobos ($n=5$) for each plate run. Faecal samples were ground to a powder using a mortar and pestle, weighed out to the protocol's recommended ≥ 0.2 g. of faecal material, avoiding any plant or partially digested food material. Samples were then diluted (1:4) in assay buffer. The kit manufacturer reported the detection limit for this assay as 45.4 pg ml^{-1} . To control for shifts in circadian rhythm for FGMC, we used those samples collected at the same time of day, specifically in the morning, under night nests, to ensure all bonobo samples were from approximately the same time point. All plates were read using a BioTek microplate reader and analysed with Gen5 software version 2.0. For the FGMC controls, 100 μl aliquots of assay buffer were divided into seven aliquots. We then spiked six of the aliquots with 100 μl of standards such that each aliquot of the sample received one of the six concentrations of standard (1000, 500, 250, 125, 62.5, 31.2 pg ml^{-1}), and one aliquot was left neat following the kit protocol to produce a standard curve. Both the spiked and neat aliquots were assayed according to kit instructions.

16s sequencing – gut microbiota composition

We used the 218 RNAlater preserved faecal samples to extract, amplify and sequence microbial DNA. Total genomic DNA was extracted from each faecal sample using the QIAamp PowerFecal DNA kit (QIAGEN) in the Ting Lab at the University of Oregon. Negative controls were included in extraction batches to test for contamination. DNA extracts were quantified using a Qubit dsDNA HS Assay Kit protocol using a Qubit 2.0 Fluorometer (Thermo Fisher Scientific). Samples containing at least 1.0 $\text{ng}/\mu\text{l}$ were sent for amplification and sequencing of the V4 hypervariable region of the bacterial 16S ribosomal RNA using 515F/806R primers at the Genomics and Cell Characterization Core Facility at the University of Oregon following previously published methods [70, 71]. Barcoded amplicons were sequenced on a 150 PE V3 run on the Illumina MiSeq platform (Illumina, San Diego, CA). The resulting sequences were demultiplexed and denoised using DADA2, and amplicon sequence variants (ASVs) were assigned using the Green Genes database. Quality filtering and assembly were done using the QIIME2 pipeline for microbial analyses [72]. The ASV table was created for samples rarified to an average sampling depth of 79,058 reads per sample. We removed samples below 315 and above 100,000 reads per sample, which accounted for 8.7% of the original dataset. We retained 202 samples for a total of 26,010,213 reads. Negative control samples were sequenced for each extraction, PCR, and library preparation. Any ASVs that appeared in these negative controls were removed from the 202 samples in the R package 'decontam' using the prevalence and frequency methods [73].

DATA ANALYSIS

We tested sex, age, whether a female had an infant, and FGMC as our predictor variables with bonobo gut microbiota composition and diversity as the response variable. We included these variables because they are associated with differences in FGMC and gut microbiota [74–76]. Statistics were run in R version 4.0.2 [77]. We included sample ID as a random effect in our models. We estimate 26–38 individuals sampled with an estimated resampling rate of 2–11 times during the data collection period based on observations of individuals and previously published estimates for Iyema [58]. We calculated two measures of alpha diversity, Shannon's and Chao's diversity indices, using the 'vegan' package [78]. We ran two-way analysis of variance (ANOVAs) against the predictor variables, against sex ($n=55$) and age ($n=59$), sex ($n=55$) and whether a female had an infant ($n=59$), and age ($n=59$) and whether a female had an infant ($n=59$). We ran linear regressions for FGMC ($n=202$) against Shannon's index and Chao's index to study alpha diversity or within individual diversity (Table 1). To examine the relationship between FGMC and bonobo gut microbiota beta diversity, we ran a permutational multivariate analysis of variance (PERMANOVA) with 999 permutations using the 'adonis2' function in the R package [79]. PERMANOVAs use the calculated beta diversity for the Jensen–Shannon distance, weighted UniFrac and unweighted UniFrac dissimilarity matrices taking the model predictors FGMC, sex, age, and whether or not a female had an infant sequentially ($n=202$) (Table 1). We used Jensen–Shannon's distance because it is useful for examining compositional differences [80]. Weighted UniFrac is a metric used to detect differences based on commonly abundant taxa [81]. At the same time, unweighted UniFrac is better at detecting differences in rare or non-abundant taxa in a community [81]. It is of note that PERMANOVAs factor in the order in which variables are entered into the model, therefore we ran this with the factors in different orders and found that the pattern of significance stayed consistent despite the order of the predictor variables. We also ran mantel tests on the log transformed FGMC values and the three beta dissimilarity matrices. We also ran abundance models on our filtered data, and ran 302 taxa using the analysis of composition of microbiomes (ANCOM) R package to test whether a member of the gut microbiota varies with high (21540–7115 ng pl^{-1}) and low concentrations (1073–7115 ng

Table 1. Predictor variables included in analysis and collection method. Sample sizes are in parentheses

| Predictor variable | Factor levels | Collection method | Lab analysis |
|--|---|---|-----------------------------------|
| FGMC | Continuous value (202) | Non-invasive faecal sample collection | ELISA assays to quantify cortisol |
| Sex | Male (24), female (31) | Behavioural observations corroborated with genetic sexing assay | Sexing assay |
| Age | Adult (48), sub-adult (4), juvenile (5), infant (2) | Behavioural observations | -- |
| Infant (whether or not a female had an infant) | Yes (10), no (49) | Behavioural observations | -- |

pl⁻¹) of FGMC ($n=202$) [82]. The cut-off was used because ANCOM requires a categorical variable of about equal sample sizes to run. All models were run with FGMC as a continuous variable except the ANCOM results, which required the FGMC values to be coded as high or low. We then subsetted out the significant ASVs and verified with general linear models with poisson distribution that those significant taxa had a linear relationship with FGMC.

RESULTS

Alpha diversity

Bonobo's faecal microbiome had similar alpha diversity regardless of FGMC levels, their sex, age or whether a female had an infant. Shannon's and Chao's indexes were not significantly related to FGMC (Fig. 1). The ANOVA results were all not significant for Shannon's and Chao's diversity indices against sex (Shannon: P -value=0.919; Chao's: P -value=0.488), age (Shannon: P -value=0.955; Chao's: P -value=0.699), and whether or not a female had an infant (Shannon: P -value=0.912; Chao's: P -value=0.521) (Fig. 2).

Bonobo's FGMC did not change due to sex, age, and whether or not a female had an infant. We found no significant interaction between FGMC and sex (Shannon: P -value=0.131; Chao's: P -value=0.510), age (Shannon: P -value=0.143; Chao's: P -value=0.459), and whether or not a female had an infant (Shannon: P -value=0.131; Chao's: P -value=0.487) for both alpha diversity metrics when run in two-way ANOVA with an interaction effect. FGMC values had mean $7529 \text{ ng pl}^{-1} \pm 266.55 \text{ ng pl}^{-1}$ and did not significantly differ by sex (P -value=0.387), age (P -value=0.17), and whether or not a female had an infant (P -value=0.144).

Beta diversity

Bonobo FGMC significantly explained a small amount of variation in beta diversity for two of the three beta diversity metrics, Jensen–Shannon's distance and weighted UniFrac. Beta diversity did not significantly relate to FGMC for unweighted UniFrac. The PERMANOVA results for FGMC showed that it explained 0.8% of the variation in beta diversity for the Jensen–Shannon distance (Table 2). The PERMANOVA for the weighted UniFrac dissimilarity matrix found FGMC explained 1.2% (Fig. 3; Table 2). The PERMANOVA results for FGMC showed that the unweighted UniFrac dissimilarity matrix did not significantly explain variation in beta diversity (Table 2).

Mantel test showed that there was a significant agreement between the FGMC Euclidian distance matrix and the Jensen–Shannon distance matrix showing that large difference in FGMC value were associated with high values of dissimilarity in Jensen–Shannon distance (observed value: 0.09, P -value=0.02) (Fig. 4). The mantel test for weighted UniFrac (P -value=0.28) and unweighted UniFrac (P -value=0.17) were not significant.

Analysis of community variance (ANCOM) results

Bonobos have 17 taxa whose abundance was explained by FGMC for the ANCOM model after Bonferroni correction out of 302 taxa tested (Table 3; Fig. S1 available with the online version of this article.). We found 15 of the significant ASVs had a positive linear relation with FGMC while two significant ASVs *RFN20* and *Butyrivibrio*, had a negative linear relation with FGMC.

DISCUSSION

We aimed to test the relationship between FGMC and the gut microbiota for bonobos and compare the results to those reported for western lowland gorillas, humans and other mammals [20, 28–30, 35]. We predicted that the bonobo gut microbiota would exhibit patterns similar to what was found in western lowland gorillas, where alpha diversity (or within individual diversity) and beta diversity (or between individual diversity) were not significantly explained by FGMC due to the similar social environment, ecology and phylogenetic relationship between western lowland gorillas and bonobos. The abundance model results for the western lowland gorillas found three taxa significantly correlated with FGMC [35].




Fig. 1. Scatterplots of FGMC against (a) Shannon's diversity and (b) Chao's diversity index were not significant.

Consistent with Vlčková *et al.* [35], our alpha diversity measures were not significant. This suggests that number of bacterial taxa within the gut microbial community stays constant. Additionally, it suggests there is stability in the number of taxa in the guts of great apes even when the gut microbiome is disrupted by stress. This is constant with several other findings outside of great apes, where despite potentially stressful habitat fragmentation, red colobus (*Procolobus rufomitratus*), black and white colobus (*Colobus guerza*) and red-tailed guenon (*Cercopithecus ascanius*) gut microbiomes remained stable [83]. This stability in alpha diversity may be a wild primate feature as other mammals like elephants [29], pangolins [28] and squirrels [27, 84] have been found to exhibit changes in alpha diversity linked to FGMC. This may point to the ability of primates to buffer stressful events due to their behavioural flexibility and sociality [84, 85].

Our beta diversity results found that FGMC significantly explains a small amount of between individual variation in the bonobo gut microbiota. The amount of variation explained in our PERMANOVA by FGMC was very small and could be due noise in the data, but because we also detected a difference with the mantel tests for the Jensen-Shannon dissimilarity metric this result is likely an actual pattern. The small amount of variation explained implies that there is only a small number of taxa whose abundance is affected by FGMC. In other primates, beta diversity has been found to significantly change with degraded habitats in howler monkeys (*Alouatta pigra*), potentially due to the stress of inhabiting and eating much lower quality food items [86]. In contrast, beta diversity for red colobus (*Procolobus rufomitratus*), black and white colobus (*Colobus guerza*), and red-tailed




Fig. 2. The predictor variables, sex (column 1), age (column 2) and infant (column 3) were all not significant for (a). Shannon's and (b). Chao's.

guenon in degraded habitats remained stable [83]. In primates, there does not seem to be a clear pattern of how stress and beta diversity relate to each other and may depend on the context that a wild primate is living in or may depend on the specific taxa found in the gut of a wild-living primate. In mammals, elephants [29] and pangolins [28], beta diversity significantly changes with a stress. Beta diversity changes in response to stress appear to be context and species specific. The fact that beta diversity was significant for bonobos and not gorillas implies that bonobos gut microbiomes may be more susceptible to stress or that bonobo gut microbiomes are home to bacteria whose abundance is more susceptible to fluctuations in stress.

There were 17 taxa in the bonobo gut microbiome whose abundance was significantly related to FGMC, for both our ANCOM and linear model results. Vlčková *et al.* [35] found only three microbial taxa were shown to be significant with FGMC in western lowland gorillas (family Anaerolineaceae; genus *Clostridium* cluster XIVb; genus *Oscillibacter*) [35]. We found members of the family Anaerolineaceae in bonobo samples, and interestingly one taxon (genus SHD-231) was significant in our differential

Table 2. PERMANOVA results for Jensen–Shannon distance, weighted UniFrac, and unweighted UniFrac against FGMC, sex, age and whether or not a female had an infant

| | <i>Df</i> | Sums of squares | Mean squares | <i>F-value</i> | <i>R</i> ² | <i>P-value</i> |
|--------------------|-----------|-----------------|--------------|----------------|-----------------------|----------------|
| Jensen-Shannon | | | | | | |
| FGMC | 1, 195 | 0.29 | 0.29 | 1.77 | 0.008 | 0.003** |
| Sex | 2, 195 | 0.32 | 0.16 | 0.97 | 0.009 | 0.54 |
| Age | 3, 195 | 0.47 | 0.15 | 0.96 | 0.014 | 0.56 |
| Infant | 1, 195 | 0.15 | 0.15 | 0.95 | 0.004 | 0.56 |
| weighted UniFrac | | | | | | |
| FGMC | 1, 195 | 0.09 | 0.09 | 2.57 | 0.012 | 0.03 * |
| Sex | 2, 195 | 0.05 | 0.02 | 0.84 | 0.008 | 0.52 |
| Age | 3, 195 | 0.06 | 0.02 | 0.63 | 0.009 | 0.72 |
| Infant | 1, 195 | 0.03 | 0.03 | 0.91 | 0.004 | 0.40 |
| unweighted UniFrac | | | | | | |
| FGMC | 1, 195 | 0.26 | 0.26 | 1.98 | 0.009 | 0.06 |
| Sex | 2, 195 | 0.21 | 0.10 | 0.77 | 0.007 | 0.65 |
| Age | 3, 195 | 0.37 | 0.12 | 0.91 | 0.01 | 0.50 |
| Infant | 1, 195 | 0.11 | 0.11 | 0.87 | 0.004 | 0.47 |




Fig. 3. .PCoA plot of weighted UniFrac dissimilarity matrix. The PERMANOVA results suggest that FGMC explained 1.27% of the variation in beta diversity.

abundance model and linear model results, similar to gorillas [35]. We did find one genus of *Clostridium* in the 201 bonobos that we sampled; however, this genus of *Clostridium* was not a part of cluster XIVb, found in western lowland gorillas to be significant [35]. We found no genus *Oscillibacter*, a genus found to be significant in gorillas, in our bonobo samples, nor did we detect any of the higher family level Oscillospiraceae. Therefore, bonobos and western lowland gorillas appear to be similar in that members of the family Anaerolineaceae may be differentially affected during periods of high stress but differ in the 16 other taxa that are differentially abundant based on FGMC in bonobos.

Other taxa that we found specific to the bonobo abundance results are two different unknown bacteria in the order Clostridiales that are thought to be linked to early life stress in mice [87]. Additionally, two members of the order Bacteroidales, including family S24-7, which has been associated with changes in circadian rhythm disruption in murine models [88]. We also found Ruminococcaceae and Mogibacteriaceae, which have been found in the human gut microbiota [89, 90]. Other notable genus level associations included *Prevotella* sp., associated with chronic inflammatory conditions [91]. These different patterns between the bonobo and western lowland gorillas suggest that there may be species-specific or temporal-specific effects of FGMC on the abundance of specific taxa in primate gut microbiota, and several of these taxa have been linked to early life stress, circadian




Fig. 4. Frequency distribution from the Mantel test showing the results of the 999 randomizations for Jensen–Shannon distance matrix and the FGMC Euclidian distance matrix. The observed value is marked with the line and black diamond.

Table 3. Taxa from ANCOM results were significantly related to FGMC after Bonferroni correction. *W* is equal to the number of times the log-ratio of a particular taxon compared to every other taxon being tested was detected to be significantly different across groups. For all the listed taxa, we accepted the alternative hypothesis that the FGMC does significantly explain the abundance of the listed taxa. The taxa are listed by kingdom, phylum, class, order, family, genus. Not all ASVs were identified to the genus or family level.

| Classification of Amplicon Sequence Variant (ASV) | w | Linear regression results (FGMC ~ ASV) |
|---|-----|---|
| Bacteria, Firmicutes, Clostridia, Clostridiales | 272 | $4.706 \times 10^{-05} \pm 9.766 \times 10^{-07}$, <i>P</i> -value = $< 2 \times 10^{-16}$ *** (Fig. S2) |
| Archaea, Euryarchaeota, Methanobacteria, Methanobacteriales, Methanobacteriaceae, <i>Methanobrevibacter</i> | 268 | $4.909 \times 10^{-05} \pm 6.995 \times 10^{-07}$, <i>P</i> -value = $< 2 \times 10^{-16}$ *** (Fig. S3) |
| Bacteria, Firmicutes, Clostridia, Clostridiales | 266 | $5.811 \times 10^{-05} \pm 1.052 \times 10^{-06}$, <i>P</i> -value = $< 2 \times 10^{-16}$ *** (Fig. S4) |
| Bacteria, Bacteroidetes, Bacteroidia, Bacteroidales | 266 | $1.223 \times 10^{-04} \pm 1.206 \times 10^{-06}$, <i>P</i> -value = $< 2 \times 10^{-16}$ *** (Fig. S5) |
| Archaea, Euryarchaeota, Thermoplasmata, E2, Methanomassiliococcaceae, <i>vadinCA11</i> | 260 | $1.107 \times 10^{-04} \pm 1.148 \times 10^{-06}$, <i>p</i> -value = $< 2 \times 10^{-16}$ *** (Fig. S6) |
| Bacteria, Proteobacteria, Betaproteobacteria, Burkholderiales | 255 | $5.640 \times 10^{-04} \pm 6.092 \times 10^{-06}$, <i>P</i> -value = $< 2 \times 10^{-16}$ *** (Fig. S7) |
| Bacteria, Bacteroidetes, Bacteroidia, Bacteroidales, RF16 | 253 | $2.829 \times 10^{-04} \pm 2.420 \times 10^{-06}$, <i>P</i> -value = $< 2 \times 10^{-16}$ *** (Fig. S8) |
| Bacteria, Bacteroidetes, Bacteroidia, Bacteroidales, Prevotellaceae, <i>Prevotella</i> | 249 | $1.615 \times 10^{-05} \pm 1.452 \times 10^{-07}$, <i>P</i> -value = $< 2 \times 10^{-16}$ *** (Fig. S9) |
| Bacteria, Chloroflexi, Anaerolineae, Anaerolineales, Anaerolinaceae, <i>SHD-231</i> | 247 | $6.985 \times 10^{-04} \pm 1.244 \times 10^{-06}$, <i>P</i> -value = $< 2 \times 10^{-16}$ *** (Fig. S10) |
| Bacteria, Bacteroidetes, Bacteroidia, Bacteroidales, Paraprevotellaceae | 247 | $2.020 \times 10^{-04} \pm 2.521 \times 10^{-06}$, <i>P</i> -value = $< 2 \times 10^{-16}$ *** (Fig. S11) |
| Bacteria, Bacteroidetes, Bacteroidia, Bacteroidales, S24-7 | 247 | $2.393 \times 10^{-04} \pm 3.147 \times 10^{-06}$, <i>P</i> -value = $< 2 \times 10^{-16}$ *** (Fig. S12) |
| Bacteria, Firmicutes, Clostridia, Clostridiales, Ruminococcaceae | 246 | $4.715 \times 10^{-05} \pm 6.250 \times 10^{-07}$, <i>P</i> -value = $< 2 \times 10^{-16}$ *** (Fig. S13) |
| Bacteria, Firmicutes, Erysipelotrichi, Erysipelotrichales, Erysipelotrichaceae, <i>RFN20</i> | 241 | $-5.490 \times 10^{-04} \pm 1.261 \times 10^{-05}$, <i>P</i> -value = $< 2 \times 10^{-16}$ *** (Fig. S14) |
| Bacteria, Verrucomicrobia, Verruco-5, WCHB1-41, RFP12 | 241 | $5.358 \times 10^{-05} \pm 3.110 \times 10^{-06}$, <i>P</i> -value = $< 2 \times 10^{-16}$ *** (Fig. S15) |
| Bacteria, Firmicutes, Clostridia, Clostridiales, Ruminococcaceae | 239 | $1.713 \times 10^{-04} \pm 1.950 \times 10^{-06}$, <i>P</i> -value = $< 2 \times 10^{-16}$ *** (Fig. S16) |
| Bacteria, Firmicutes, Clostridia, Clostridiales, Mogibacteriaceae | 235 | $2.899 \times 10^{-04} \pm 5.295 \times 10^{-06}$, <i>P</i> -value = $< 2 \times 10^{-16}$ *** (Fig. S17) |
| Bacteria, Firmicutes, Clostridia, Clostridiales, Lachnospiraceae, <i>Butyrivibrio</i> | 229 | $-1.289 \times 10^{-04} \pm 2.256 \times 10^{-06}$, <i>P</i> -value = $< 2 \times 10^{-16}$ *** (Fig. S18) |

rhythm stress and chronic inflammation [26, 87–90]. However, it does appear that the family Anaerolinaceae may be particularly affected by stress in great apes.

In other mammals, like squirrels, pangolins, elephants and rhinos, stress was associated with changes, in specific taxa [27–30, 92]. Additionally, specific taxa changed in abundance due to stress or stressful events [27, 28, 30]. The specific taxa are differentially expressed in elephants (Planococcaceae, Clostridiaceae, Spirochaetaceae and Bacteroidia) and rhinos (Aerococcaceae, Atopostipes, Carnobacteriaceae and Solobacterium) were not found to be related explicitly to FGMC in bonobos [29]. However, there may be high-order similarities between bonobos and other mammals in the specific taxa that are differential abundant due to stress. Clostridiaceae is a family belonging to the class Clostridia that was found to be differentially expressed in elephants [29]. While we did not find this specific family differential expressed in bonobos, we found six Clostridia members to be differentially expressed in bonobos. This result may indicate that stress may affect specific taxa in the gut microbiota across mammalian lineages, including humans. In humans, the proposed mechanism for a host and its associated microbes to communicate is through hormones [14, 19]. We did not see an overall decrease in diversity as has been seen in humans, but we did see certain taxa whose abundance seems to be differentially affected by stress [10, 11, 89, 91]. Indicating that there may be similarities between humans and one of their closest evolutionary relatives, bonobos.

There are several differences in methods that could be influencing stress and the gut microbiota between bonobos and gorillas. One of the significant differences between our study and the gorilla study is the sample size. Our larger sample size may

be why we picked up the small effect of FGMC on beta diversity, but more studies across wild-living primates examining stress and the gut microbiota will help to elucidate how sample size plays into picking up relationships between proximate measures of stress and the gut microbiota. Other differences include, potentially different FGMC measures and differences in the method of sample preservation for FGMC analysis [35, 93]. There are also species-specific patterns in the production, metabolism and excretion of FGMCs [74, 93, 94]. Therefore, comparing FGMC values between bonobos and gorillas must be done with extreme care. Since we are not directly comparing our FGMC values to those obtained for the western lowland gorillas the broad patterns from comparing those FGMCs to the gut microbiota may be species specific. Additionally, the gut microbiota may be responding to FGMCs in non-linear ways, and there may be more nuanced changes to consider when comparing stress and the NHP gut microbiota.

Other limitations include the metabolism of FGMCs can also be dependent on sex and time of day [95]. At the same time, we attempted to control for this variation by only selecting samples collected around the same time of day and including sex and age as factors in our analyses. This method of controlling for time of day is like the western lowland gorilla paper, where morning faecal samples were analysed for the unhabituated groups [35]. In addition to variation in FGMCs, there are several other factors that influence the composition and diversity of gut microbiota among NHP.

There are other variables that we did not examine in this paper that have been thought to influence the composition and diversity of the gut microbiota in NHP. Disease status could be influencing the gut microbiota [13, 96]. Rank and other social factors like rates of affiliation and aggression both within and between communities could also be affecting the gut microbiota [71, 96, 97]. Diet and seasonality can also be significant factors in changing how nutritionally stressed an individual is and can directly affect the gut microbiota composition [98–100]. We aim to examine these factors in future analyses.

Future directions for this work include adding metagenomic sequencing and metabolomic data to our dataset to incorporate more functional results. Additionally, we aim to incorporate diet, food availability and social variables in future analyses of the bonobo gut microbiota. Compared to humans, in bonobos, beta diversity and some taxa change in abundance instead of broadly changing gut microbial diversity in response to stress, but we did find similarities in bonobos and gorillas in the family Anaerolinaceae. Thus, future studies should examine how Anaerolinaceae changes in response to stress in other great apes like humans, chimpanzees and orangutans. Incorporating FGMC and gut microbiota data can provide a more robust understanding of how stress impacts the gut microbiota of primates.

Funding information

We would like to acknowledge the many funding agencies that have supported this research. The 2017 field season was funded thanks to a grant from the Leakey Foundation to MLW, grants, and fellowships from Northern Kentucky University to MLW, a Professional Development grant from a National Geographic grant to CMB (WW-099ER-17), and a Global Oregon Award to CMB. Grants and fellowships funded the lab work from Northern Kentucky University to MLW, a University of Oregon Gary Smith Summer Development grant, a University of Oregon Department of Anthropology Research grant to AJH, and a University of Oregon Faculty Research Award to FJW.

Acknowledgements

We want to thank the following people without whom this work would not be possible: Jef Dupain, Hugues Akpona, and the African Wildlife Foundation, Dipon Bomposo, Beken Bompoma, Teddy Bofaso, Christian Djambo, Mathieu Esaola, Bellevie Iyambe, Augustin Lofili, Isaac Lokoli, Gedeon Lokofu, Thomas Lokuli, Joel Bontambe, Abdulay Bokela, Papa Siri, Paco Bertolani, Ian Takaoka, and many others who generously provided both logistical support and friendship in the field. Additionally, we want to thank Diana Christie and the Molecular Anthropology laboratory group members for their helpful discussions through all stages of this research.

Author contributions

A.J.H., K.J.B., M.L.W., C.M.B., F.J.W., J.J.S., N.T. Preparation, creation and/or presentation of the published work by those from the original research group, specifically critical review, commentary or revision – including pre- or post-publication stages. A.J.H. Preparation, creation and/or presentation of the published work, specifically visualization/data presentation. F.J.W. Oversight and leadership responsibility for the research activity planning and execution, including mentorship external to the core team. F.J.W., M.L.W., N.T. Management and coordination responsibility for the research activity planning and execution. A.J.H., F.J.W., M.L.W., C.M.B. Acquisition of the financial support for the project leading to this publication.

Conflicts of interest

The authors have no conflicts of interest to declare.

Ethical statement

This research was completed with the approval of the University of Oregon's Institutional Animal Care and Use Committee (AUP-17-10), abided by the American Society of Primatologist's Principles for the Ethical Treatment of Human Primates, and was approved by the L'Institut Congolais pour la Conservation de la Nature (ICCN) (permit no. 3058) the department within the Democratic Republic of the Congo that oversees the Lomako-Yokokala Faunal Reserve where the Iyema field site is located. Permissions were obtained through the African Wildlife Foundation Mission Order (No. 033/2017).

References

1. Allaband C, McDonald D, Vázquez-Baeza Y, Minich JJ, Tripathi A, et al. Microbiome 101: studying, analyzing, and interpreting gut microbiome data for clinicians. *Clin Gastroenterol Hepatol* 2019;17:218–230.
2. Amato KR. Incorporating the gut microbiota into models of human and non-human primate ecology and evolution: gut microbiota influences host nutrition, health, and behavior. *Am J Phys Anthropol* 2016;159.

3. Dantas G, Sommer MOA, Degnan PH, Goodman AL. Experimental approaches for defining functional roles of microbes in the human gut. *Annu Rev Microbiol* 2013;67:459–475.
4. Barlow GM, Yu A, Mathur R. Role of the gut microbiome in obesity and diabetes mellitus. *Nutr Clin Pract* 2015;30:787–797.
5. Chattopadhyay I, Dhar R, Pethusamy K, Seethy A, Srivastava T, et al. Exploring the role of gut microbiome in colon cancer. *Appl Biochem Biotechnol* 2021;193:1780–1799.
6. Franzosa EA, Sirota-Madi A, Avila-Pacheco J, Fornelos N, Haiser HJ, et al. Gut microbiome structure and metabolic activity in inflammatory bowel disease. *Nat Microbiol* 2019;4:293–305.
7. Kim YS, Milner JA. Dietary modulation of colon cancer risk. *J Nutr* 2007;137:2576S–2579S.
8. Turnbaugh PJ, Hamady M, Yatsunenko T, Cantarel BL, Duncan A, et al. A core gut microbiome in obese and lean twins. *Nature* 2009;457:480–484.
9. Konturek PC, Brzozowski T, Konturek SJ. Stress and the gut: pathophysiology, clinical consequences, diagnostic approach and treatment options. *J Physiol Pharmacol* 2011;62:591–599.
10. Bharwani A, Mian MF, Foster JA, Surette MG, Bienenstock J, et al. Structural & functional consequences of chronic Psychosocial stress on the microbiome & host. *Psychoneuroendocrinology* 2016;63:217–227.
11. Bhatia V, Tandon RK. Stress and the gastrointestinal tract. *J Gastroenterol Hepatol* 2005;20:332–339.
12. Bailey MT. The effects of psychological stressors on the intestinal microbiota. *Biosci Microflora* 2009;28:125–134. ;
13. Moloney RD, Desbonnet L, Clarke G, Dinan TG, Cryan JF. The microbiome: stress, health and disease. *Mamm Genome* 2014;25:49–74.
14. Tetel MJ, de Vries GJ, Melcangi RC, Panzica G, O'Mahony SM. Steroids, stress and the gut microbiome-brain axis. *J Neuroendocrinol* 2018;30.
15. Söderholm JD, Perdue MH. II. Stress and intestinal barrier function. *Am J Physiol Gastrointest Liver Physiol* 2001;280:G7–G13.
16. Collins SM. Stress and the gastrointestinal tract IV. Modulation of intestinal inflammation by stress: basic mechanisms and clinical relevance. *Am J Physiol Gastrointest Liver Physiol* 2001;280:G315–8.
17. Bailey MT, Coe CL. Maternal separation disrupts the integrity of the intestinal microflora in infant rhesus monkeys. *Dev Psychobiol* 1999;35:146–155.
18. Keay JM, Singh J, Gaunt MC, Kaur T. Fecal glucocorticoids and their metabolites as indicators of stress in various mammalian species: a literature review. *J Zoo Wildl Med* 2006;37:234–244.
19. Sandrini S, Aldriwesh M, Alruways M, Freestone P. Microbial endocrinology: host-bacteria communication within the gut microbiome. *J Endocrinol* 2015;225:R21–34.
20. Foster JA, Rinaman L, Cryan JF. Stress & the gut-brain axis: regulation by the microbiome. *Neurobiol Stress* 2017;7:124–136.
21. Moser V. Graduate research projects. *The human microbiome: the brain-gut axis and its role in immunity* 2014:7.
22. Crumeyrolle-Arias M, Jaglin M, Bruneau A, Vancassel S, Cardona A, et al. Absence of the gut microbiota enhances anxiety-like behavior and neuroendocrine response to acute stress in rats. *Psychoneuroendocrinology* 2014;42:207–217.
23. Hata T, Miyata N, Takakura S, Yoshihara K, Asano Y, et al. The gut microbiome derived from Anorexia Nervosa patients impairs weight gain and behavioral performance in mice. *SSRN Journal* 2018;3155625.
24. Li J, Sung CYJ, Lee N, Ni Y, Pihlajamäki J, et al. Probiotics modulated gut microbiota suppresses hepatocellular carcinoma growth in mice. *Proc Natl Acad Sci U S A* 2016;113:E1306–15.
25. Bravo JA, Forsythe P, Chew MV, Escaravage E, Savignac HM, et al. Ingestion of *Lactobacillus* strain regulates emotional behavior and central GABA receptor expression in a mouse via the vagus nerve. *Proc Natl Acad Sci U S A* 2011;108:16050–16055.
26. Bharwani A, Mian MF, Foster JA, Surette MG, Bienenstock J, et al. Structural & functional consequences of chronic psychosocial stress on the microbiome & host. *Psychoneuroendocrinology* 2016;63:217–227.
27. Stothart MR, Palme R, Newman AEM. It's what's on the inside that counts: stress physiology and the bacterial microbiome of a wild urban mammal. *Proc Biol Sci* 2019;286:20192111.
28. Yan D, Hu D, Li K, Li B, Zeng X, et al. Effects of chronic stress on the Fecal Microbiome of Malayan Pangolins (*Manis javanica*) rescued from the illegal wildlife trade. *Curr Microbiol* 2021;78:1017–1025.
29. Moustafa MAM, Chel HM, Thu MJ, Bawm S, Htun LL, et al. Anthropogenic interferences lead to gut microbiome dysbiosis in Asian elephants and may alter adaptation processes to surrounding environments. *Sci Rep* 2021;11:741.
30. Antwis RE, Edwards KL, Unwin B, Walker SL, Shultz S. Rare gut microbiota associated with breeding success, hormone metabolites and ovarian cycle phase in the critically endangered eastern black rhino. *Microbiome* 2019;7:27.
31. Surbeck M, Deschner T, Weltring A, Hohmann G. Social correlates of variation in urinary cortisol in wild male bonobos (*Pan paniscus*). *Horm Behav* 2012;62:27–35.
32. Young T, Isbell L. Ecological models of female social relationships in primates: similarities, disparities, and some directions for future clarity. *Behav* 2002;139:177–202.
33. Amato KR. Incorporating the gut microbiota into models of human and non-human primate ecology and evolution. *Am J Phys Anthropol* 2016;159:S196–215.
34. Bailey MT. The contributing role of the intestinal microbiota in stressor-induced increases in susceptibility to enteric infection and systemic immunomodulation. *Horm Behav* 2012;62:286–294.
35. Vlčková K, Shutt-Phillips K, Heistermann M, Pafčo B, Petrželková KJ, et al. Impact of stress on the gut microbiome of free-ranging western lowland gorillas. *Microbiology* 2018;164:40–44.
36. McGrew WC, Marchant LF, Nishida T, Goodall J, Itani J. *Great Ape Societies*. Cambridge University Press, 1996.
37. Tan J, Ariely D, Hare B. Bonobos respond prosocially toward members of other groups. *Sci Rep* 2017;7:14733.
38. White FJ. *Pan paniscus* 1973 to 1996: twenty-three years of field research. *Evol Anthropol* 1996;5:11–17.
39. Gruber T, Clay Z. A comparison between Bonobos and Chimpanzees: a review and update. *Evol Anthropol* 2016;25:239–252.
40. Forcina G, Vallet D, Le Gouar PJ, Bernardo-Madrid R, Illera G, et al. From groups to communities in western lowland gorillas. *Proc Biol Sci* 2019;286:20182019.
41. Parnell RJ. Group size and structure in western lowland gorillas (*Gorilla gorilla gorilla*) at Mbeli Bai, Republic of Congo. *Am J Primatol* 2002;56:193–206.
42. Harcourt AH, Stewart KJ. Gorilla society: what we know and don't know. *Evol Anthropol* 2007;16:147–158. ;
43. Kyrou I, Tsigos C. Stress hormones: physiological stress and regulation of metabolism. *Curr Opin Pharmacol* 2009;9:787–793.
44. Thau L, Gandhi J, Sharma S. *Physiology, Cortisol*. StatPearls Publishing, 2021.
45. Tkaczynski PJ, Behringer V, Ackermann CY, Fedurek P, Fruth B, et al. Patterns of urinary cortisol levels during ontogeny appear population specific rather than species specific in wild chimpanzees and bonobos. *J Hum Evol* 2020;147:102869.
46. Fourie NH, Brown JL, Jolly CJ, Phillips-Conroy JE, Rogers J, et al. Sources of variation in hair cortisol in wild and captive non-human primates. *Zoology (Jena)* 2016;119:119–125.
47. Strier KB, Ziegler TE, Wittwer DJ. Seasonal and social correlates of fecal testosterone and cortisol levels in wild male muriquis (*Brachyteles arachnoides*). *Horm Behav* 1999;35:125–134.
48. Hohmann G, Mundry R, Deschner T. The relationship between socio-sexual behavior and salivary cortisol in bonobos: tests of the tension regulation hypothesis. *Am J Primatol* 2009;71:223–232.

49. Mason HL, Myers CS, Kendall EC. Chemical studies of the suprarenal cortex II. the identification of a substance which possesses the qualitative action of cortin; its conversion into a diketone closely related to androstenedione. *J Biol Chem* 1936;116:267–276.
50. Mason HL, Hoehn WM, Kendall EC. Chemical studies of the suprarenal cortex. *J Biol Chem* 1938;124:459–474. ;
51. Millspaugh JJ, Washburn BE. Use of fecal glucocorticoid metabolite measures in conservation biology research: considerations for application and interpretation. *Gen Comp Endocrinol* 2004;138:189–199.
52. Davenport ER, Sanders JG, Song SJ, Amato KR, Clark AG, et al. The human microbiome in evolution. *BMC Biol* 2017;15:127.
53. Cheng L, Lucchesi S, Mundry R, Samuni L, Deschner T, et al. Variation in aggression rates and urinary cortisol levels indicates intergroup competition in wild bonobos. *Horm Behav* 2021;128:104914.
54. Cobden AK. *Party Animals: Food, Sociality and Stress in Wild Bonobos (Pan paniscus) of Iyema, Lomako Forest, Democratic Republic of Congo, PhD Thesis*. Emory University, 2014.
55. Malenky RK, Wrangham RW. A quantitative comparison of terrestrial herbaceous food consumption by *Pan paniscus* in the Lomako Forest, Zaire, and *Pan troglodytes* in the Kibale Forest, Uganda. *Am J Primatol* 1994;32:1–12. ;
56. Wrangham RW, White FJ. Feeding competition and patch size in the chimpanzee species *Pan paniscus* and *Pan troglodytes*. *Behav* 1988;105:148–164.
57. Dupain J, Van Krunkelsven E, Van Elsacker L, Verheyen RF. Current status of the bonobo (*Pan paniscus*) in the proposed Lomako Reserve (Democratic Republic of Congo). *Biol Conserv* 2000;94:265–272.
58. Brand CM, White FJ, Wakefield ML, Waller MT, Ruiz-Lopez MJ, et al. Initiation of genetic demographic monitoring of bonobos (*Pan paniscus*) at Iyema, Lomako forest, DRC. *Primate Conserv News J IUCNSSC Primate Spec Group* 2016;30:103.
59. Prado-Martinez J, Sudmant PH, Kidd JM, Li H, Kelley JL, et al. Great ape genetic diversity and population history. *Nature* 2013;499:471–475.
60. Langergraber KE, Prüfer K, Rowney C, Boesch C, Crockford C, et al. Generation times in wild chimpanzees and gorillas suggest earlier divergence times in great ape and human evolution. *Proc Natl Acad Sci U S A* 2012;109:15716–15721.
61. Blekhman R, Tang K, Archie EA, Barreiro LB, Johnson ZP, et al. Common methods for fecal sample storage in field studies yield consistent signatures of individual identity in microbiome sequencing data. *Sci Rep* 2016;6:31519.
62. Hayakawa T, Sawada A, Tanabe AS, Fukuda S, Kishida T, et al. Improving the standards for gut microbiome analysis of fecal samples: insights from the field biology of Japanese macaques on Yakushima Island. *Primates* 2018;59:423–436.
63. Vlčková K, Mrázek J, Kopečný J, Petrželková KJ. Evaluation of different storage methods to characterize the fecal bacterial communities of captive western lowland gorillas (*Gorilla gorilla gorilla*). *J Microbiol Methods* 2012;91:45–51.
64. Hale VL, Tan CL, Knight R, Amato KR. Effect of preservation method on spider monkey (*Ateles geoffroyi*) fecal microbiota over 8 weeks. *J Microbiol Methods* 2015;113:16–26.
65. Gorzelak MA, Gill SK, Tasnim N, Ahmadi-Vand Z, Jay M, et al. Methods for improving human gut microbiome data by reducing variability through sample processing and storage of stool. *PLoS ONE* 2015;10:e0134802.
66. Choo JM, Leong LEX, Rogers GB. Sample storage conditions significantly influence faecal microbiome profiles. *Sci Rep* 2015;5:16350.
67. Menke S, Gillingham MAF, Wilhelm K, Sommer S. Home-made cost effective preservation buffer is a better alternative to commercial preservation methods for microbiome research. *Front Microbiol* 2017;8:102.
68. Midttveit E. Measuring stress in captive bonobos: a look to the past and future to improve methods; 2016 Oct. <https://scholarbank.uoregon.edu/xmlui/handle/1794/20464> [accessed 15 January 2021].
69. Brand CM, Boose KJ, Squires EC, Marchant LF, White FJ, et al. Hair plucking, stress, and urinary cortisol among captive bonobos (*Pan paniscus*). *Zoo Biol* 2016;35:415–422.
70. Goodfellow CK, Whitney T, Christie DM, Sicotte P, Wikberg EC, et al. Divergence in gut microbial communities mirrors a social group fission event in a black-and-white colobus monkey (*Colobus vellerosus*). *Am J Primatol* 2019;81:e22966.
71. Wikberg EC, Christie D, Sicotte P, Ting N. Interactions between social groups of colobus monkeys (*Colobus vellerosus*) explain similarities in their gut microbiomes. *Anim Behav* 2020;163:17–31.
72. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, et al. Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nat Biotechnol* 2019;37:852–857.
73. Davis NM, Proctor DM, Holmes SP, Relman DA, Callahan BJ. Simple statistical identification and removal of contaminant sequences in marker-gene and metagenomics data. *Microbiome* 2018;6:226.
74. Colby HD, Kitay JI. Sex and substrate effects on hepatic corticosteroid metabolism in the rat. *Endocrinology* 1972;90:473–478.
75. Adriansjach J, Baum ST, Lefkowitz EJ, Van Der Pol WJ, Buford TW, et al. Age-related differences in the gut microbiome of rhesus macaques. *J Gerontol A Biol Sci Med Sci* 2020;75:1293–1298.
76. Aatsinki A-K, Keskitalo A, Laitinen V, Munukka E, Uusitupa H-M, et al. Maternal prenatal psychological distress and hair cortisol levels associate with infant fecal microbiota composition at 2.5 months of age. *Psychoneuroendocrinology* 2020;119:104754.
77. The R Development Core Team. R: A language and environment for statistical computing. 2013.
78. Oksanen J, Guillaume FB, Roeland K, Peter RM, Gavin LS, et al. *Package Vegan: Community ecology package, version 2.0 10*. Cran, 2013.
79. Anderson MJ. Permutational multivariate analysis of variance (PERMANOVA). *Wiley Statsref Stat Ref Online* 2014:1–15.
80. Menéndez ML, Pardo JA, Pardo L, Pardo MC. The Jensen-Shannon divergence. *J Franklin Inst* 1997;334:307–318.
81. Lozupone C, Knight R. UniFrac: a new phylogenetic method for comparing microbial communities. *Appl Environ Microbiol* 2005;71:8228–8235.
82. Mandal S, Van Treuren W, White RA, Eggesbø M, Knight R, et al. Analysis of composition of microbiomes: a novel method for studying microbial composition. *Microb Ecol Health Dis* 2015;26:27663.
83. McCord AI, Chapman CA, Weny G, Tumukunde A, Hyeroba D, et al. Fecal microbiomes of non-human primates in Western Uganda reveal species-specific communities largely resistant to habitat perturbation. *Am J Primatol* 2014;76:347–354.
84. Young C, Majolo B, Heistermann M, Schülke O, Ostner J. Responses to social and environmental stress are attenuated by strong male bonds in wild macaques. *Proc Natl Acad Sci U S A* 2014;111:18195–18200.
85. Cheney DL, Seyfarth RM. Stress and coping mechanisms in female primates. In: *Advances in the Study of Behavior*, vol. 39. Academic Press, 2009. pp. 1–44.
86. Amato KR, Yeoman CJ, Kent A, Righini N, Carbonero F, et al. Habitat degradation impacts black howler monkey (*Alouatta pigra*) gastrointestinal microbiomes. *ISME J* 2013;7:1344–1353.
87. Usui N, Matsuzaki H, Shimada S. Characterization of early life stress-affected gut microbiota. *Brain Sci* 2021;11:913.
88. Kim Y-M, Snijders AM, Brislaw CJ, Stratton KG, Zink EM, et al. Light-stress influences the composition of the murine gut microbiome, memory function, and plasma metabolome. *Front Mol Biosci* 2019;6:108.

89. Oki K, Toyama M, Banno T, Chonan O, Benno Y, *et al.* Comprehensive analysis of the fecal microbiota of healthy Japanese adults reveals a new bacterial lineage associated with a phenotype characterized by a high frequency of bowel movements and a lean body type. *BMC Microbiol* 2016;16:284.
90. Rajilić-Stojanović M, de Vos WM. The first 1000 cultured species of the human gastrointestinal microbiota. *FEMS Microbiol Rev* 2014;38:996–1047.
91. Ley RE. Prevotella in the gut: choose carefully. *Nat Rev Gastroenterol Hepatol* 2016;13:69–70.
92. Stothart MR, Bobbie CB, Schulte-Hostedde AI, Boonstra R, Palme R, *et al.* Stress and the microbiome: linking glucocorticoids to bacterial community dynamics in wild red squirrels. *Biol Lett* 2016;12:20150875.
93. Shutt K, Setchell JM, Heistermann M. Non-invasive monitoring of physiological stress in the Western lowland gorilla (*Gorilla gorilla gorilla*): validation of a fecal glucocorticoid assay and methods for practical application in the field. *Gen Comp Endocrinol* 2012;179:167–177.
94. Wheeler BC, Tiddi B, Kalbitzer U, Visalberghi E, Heistermann M. Methodological considerations in the analysis of Fecal Glucocorticoid metabolites in Tufted Capuchins (*Cebus apella*). *Int J Primatol* 2013;34:879–898.
95. Touma C, Sachser N, Möstl E, Palme R. Effects of sex and time of day on metabolism and excretion of corticosterone in urine and feces of mice. *Gen Comp Endocrinol* 2003;130:267–278.
96. Bennett G, Malone M, Sauther ML, Cuzzo FP, White B, *et al.* Host age, social group, and habitat type influence the gut microbiota of wild ring-tailed lemurs (*Lemur catta*). *Am J Primatol* 2016;78:883–892.
97. Tung J, Barreiro LB, Burns MB, Grenier J-C, Lynch J, *et al.* Social networks predict gut microbiome composition in wild baboons. *Elife* 2015;4.
98. Amato KR, Leigh SR, Kent A, Mackie RI, Yeoman CJ, *et al.* The gut microbiota appears to compensate for seasonal diet variation in the wild black howler monkey (*Alouatta pigra*). *Microb Ecol* 2015;69:434–443.
99. Bruorton MR, Davis CL, Perrin MR. Gut microflora of vervet and samango monkeys in relation to diet. *Appl Environ Microbiol* 1991;57:573–578.
100. Baniel A, Amato KR, Beehner JC, Bergman TJ, Mercer A, *et al.* Seasonal shifts in the gut microbiome indicate plastic responses to diet in wild geladas. *Microbiome* 2021;9:26.

Edited by: M. Welch and W. van Schaik

Five reasons to publish your next article with a Microbiology Society journal

1. When you submit to our journals, you are supporting Society activities for your community.
2. Experience a fair, transparent process and critical, constructive review.
3. If you are at a Publish and Read institution, you'll enjoy the benefits of Open Access across our journal portfolio.
4. Author feedback says our Editors are 'thorough and fair' and 'patient and caring'.
5. Increase your reach and impact and share your research more widely.

Find out more and submit your article at microbiologyresearch.org.